


Fuzzy Type Theory for Opinion Dynamics

Shreya Arya, Greta Coraglia, Paige North,
Sean O'Connor, Hans Riess, Ana Tenório

ACT

2022

OUR GROUP



Left to right: Greta, Shreya, Sean, Ana, Paige, Hans

FUZZY TYPE THEORY FOR OPINION DYNAMICS

The idea

The math

MODELING OPINIONS

Want to model "proof-relevant opinions"

We need

- ▶ types as opinions
- ▶ terms as proofs of/reasons for opinions
- ▶ fuzzy logic as confidence/certainty/strength of opinions

TYPE THEORIES AND FUZZY LOGIC

Enriching over a different monoidal category gives us a different type theory/logic

	<i>binary</i>	<i>fuzzy</i>
<i>propositions</i>	$\{0, 1\}$	$[0, 1]$
<i>types</i>	Set	$\Sigma_{S:\mathbf{Set}} S \rightarrow [0, 1]$

CATEGORIES AND TYPE THEORIES

$$\textit{Type theories} \rightleftarrows \textit{Categories}$$

Given a type theory we can obtain a category where:

- ▶ the objects are contexts Γ
- ▶ the morphisms are (lists of) terms

CATEGORICAL SEMANTICS

types in context a class of maps (*projections*) $\Gamma \vdash A$ type $\Gamma.A \xrightarrow{p_A} \Gamma$

terms

sections of projections

 $\Gamma \vdash a : A$ $\Gamma.A \xleftarrow[a]{p_A} \Gamma$

substitution

pullback along projections

...

...

ENRICHED CATEGORIES AND FUZZY TYPES

Our strategy: **enrich the categories, read the type theory!**

Call $V = \Sigma_{S:\mathbf{Set}} S \rightarrow [0, 1]$ the category whose

- ▶ objects are pairs $(S, | - |_S)$ with S a set and $| - |_S : S \rightarrow [0, 1]$ a function, called *valuation*
- ▶ morphisms $f : (S, | - |_S) \rightarrow (T, | - |_T)$ are order-preserving functions between S and T

$$\text{Fuzzy type theories} \rightleftarrows V\text{-Categories}$$

INTUITION

a V -category \mathcal{C}	an agent in the system
a context	a set of beliefs
a type (in context)	a belief (and its premises)
a term of type A	a proof of the belief A

- ▶ we want definite beliefs \Rightarrow non-fuzzy types
- ▶ but their reasons might be subject to uncertainty \Rightarrow fuzzy terms

FUZZY TYPE THEORY FOR OPINION DYNAMICS

The idea

The math

PROJECTIONS AND SECTIONS

Axiom: Types are not fuzzy

For all A , $|p_A|_{\text{hom}(\Gamma.A, \Gamma)} = 1$.

Normally, terms are sections of projections, but

$$\Gamma \xrightarrow{s} \Gamma.A \xrightarrow{p_A} \Gamma$$

id (curved arrow from Γ to Γ above the straight arrows)

$$|id| = 1 \implies |p_A| \cdot |s| = 1 \implies |p_A| = |s| = 1$$

This is too much of a restriction for us!

α -SECTIONS

Definition: α -sections

We say s is a α -section of p if $p \circ s = id$ as functions and $|p| \cdot |s| \geq \alpha$

Denoted $\Gamma \vdash s :_{\alpha} A$

and we have $\frac{\Gamma \vdash s :_{\alpha} A}{\Gamma \vdash s :_{\beta} A}$ for all $\beta \leq \alpha$

SUBSTITUTION AND PULLBACKS

Classically, substitution is performed as pullback along projections. Problem is that in the enriched case we need to consider weighted pullbacks!

Weighted pullbacks are a special case of weighted limits, which replace limits in enriched settings.

This can be used to determine the universal property of weighted pullbacks in V -categories

WEIGHTED PULLBACKS

Consider a pullback in **Set**-categories

$$\begin{array}{ccc} A \times_C B & \longrightarrow & B \\ \downarrow & & \downarrow \\ A & \longrightarrow & C \end{array}$$

What in **Set**-categories is the bijection

$$\mathrm{hom}(Z, A \times_C B) \cong \mathrm{hom}(Z, A) \times_{\mathrm{hom}(Z, C)} \mathrm{hom}(Z, B)$$

can be viewed as a weighted pullback in which


$$\mathrm{hom}(Z, A \times_C B) \cong \mathrm{hom}(Z, A)^{\mathbb{1}} \times_{\mathrm{hom}(Z, C)^{\mathbb{1}}} \mathrm{hom}(Z, B)^{\mathbb{1}}$$

With this perspective, we say that a regular pullback is $(\mathbb{1}, \mathbb{1}, \mathbb{1})$ -weighted, with $\mathbb{1} = \{*\}$


FUZZY SUBSTITUTION I

We need to find reasonable weights!

- ▶ $(\mathbb{1}, \mathbb{1}, \mathbb{1})$ with $\mathbb{1}$ the terminal object doesn't work with fuzzy terms
- ▶ We can denote $\mathbb{1}_x = (\{*\}, \text{const}(x))$ to use as our weights
- ▶ $(\mathbb{1}_{\text{val}(-)}, \mathbb{1}, \mathbb{1}_{\text{val}(-)})$




SOMETHING WEIRD



But the top-left $1.A$ is obtained by pullback along a map of value α , so it isn't the same object at $1.A$.

RESOLUTION




We can denote the top-left $1.A$ as $1.A_\alpha$ and we can read

$$\frac{\vdash s :_\alpha A}{\vdash t :_1 A_\alpha}$$

as "Given a proof of A with confidence α , we can prove with confidence 1 that we can prove A with confidence α ".

FUZZY SUBSTITUTION II

$$\frac{\Gamma, x : A, \Delta \vdash t :_{\beta} B \quad \Gamma \vdash u :_{\alpha} A}{\Gamma, \Delta[u/x] \vdash t[u/x] :_{\beta} B[u/x]_{\alpha}}$$



VALIDITY

Theorem

Such a V -category satisfies (a fuzzy version of) all structural rules of (not-yet-dependent) type theory.

Therefore we have categories to encode the logical system of the agents in our system.

THE DYNAMIC

- ▶ The work of Jakob Hansen and Robert Ghrist uses a cellular sheaf $F : Inc(G) \rightarrow Vect$ to study opinion dynamics
- ▶ The work of Hans Riess and Robert Ghrist studies cellular sheaves of the form $F : Inc(G) \rightarrow Lattices$
- ▶ We want to explore $F : Inc(G) \rightarrow V-Cat$

FUTURE WORK

- ▶ Give an enriched categorical interpretation for the dependent fuzzy types (and address definitional equality);
- ▶ Replace $[0,1]$ by any ordered monoid M ;
- ▶ Explore the dynamic side

REFERENCES I



GHRIST, R., AND RIESS, H.
Cellular sheaves of lattices and the Tarski laplacian.
Homology, Homotopy and Applications 24, 1 (2022), 325–345.



HANSEN, J., AND GHRIST, R.
Opinion dynamics on discourse sheaves, 2020.



HOFMANN, M.
Syntax and semantics of dependent types.
In *Extensional Constructs in Intensional Type Theory*. Springer, 1997, pp. 13–54.

Thank you!